



Europäisches  
Patentamt

European  
Patent Office

Office européen  
des brevets

PCT/EP 03/07084  
Rec'd PCT/EP 29 DEC 2004 #1

REC'D 26 AUG 2003

WIPO

PCT

Bescheinigung

Certificate

Attestation

Die angehefteten Unterla-  
gen stimmen mit der  
ursprünglich eingereichten  
Fassung der auf dem näch-  
sten Blatt bezeichneten  
europäischen Patentanmel-  
dung überein.

The attached documents  
are exact copies of the  
European patent application  
described on the following  
page, as originally filed.

Les documents fixés à  
cette attestation sont  
conformes à la version  
initialement déposée de  
la demande de brevet  
européen spécifiée à la  
page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

02015344.1

**PRIORITY  
DOCUMENT**  
SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts;  
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets  
p.o.

R C van Dijk





Anmeldung Nr:  
Application no.: 02015344.1  
Demande no:

Anmeldetag:  
Date of filing: 10.07.02  
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

BASF AKTIENGESELLSCHAFT

67056 Ludwigshafen  
ALLEMAGNE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:  
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.  
If no title is shown please refer to the description.  
Si aucun titre n'est indiqué se référer à la description.)

Use of a gene for increasing the oil content in plants

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)  
revendiquée(s)

Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/  
Classification internationale des brevets:

C12N15/00

Am Anmeldetag benannte Vertragstaaten/Contracting states designated at date of  
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR IE IT LI LU MC NL PT SE SK TR



Use of a gene for increasing the oil content in plants

Description

5

The invention relates to the use of a gene that when expressed will increase the total amount of oil (i.e. triacylglycerols - TAG) that is produced in transgenic organisms.

10 More specifically this invention describes the identification of a gene encoding a TAG synthesis enhancing protein (TEP).

In a first embodiment, this invention is directed to the TEP protein comprising an amino acid sequence as set forth in  
 15 SEQ ID NO: 2 or a functional fragment, derivative, variant, or orthologue thereof.

The present invention further includes the nucleotide sequence as set forth in SEQ ID NO: 1, as well as portions of the genomic  
 20 sequence, the cDNA sequence, allelic variants, synthetic variants and mutants thereof. This includes sequences that are to be used as probes, vectors for transformation or cloning intermediates.

SEQ ID NO: 2 is the deduced amino acid sequence from the open  
 25 reading frame YJR098c in SEQ ID NO: 1.

Another aspect of the present invention relates to those polypeptides, which have at least 60% identity to SEQ ID NO: 2.

30 The invention furthermore relates to expression constructs for expressing yeast TEP in plants, preferably in plant seeds, transgenic plants expressing yeast TEP, and to the use of said transgenic plants for the production of food, feeds, seed, pharmaceuticals or fine chemical, in particular for  
 35 the production of oils.

In oil crops like rape, sunflower, oil palm etc., the oil (i.e. triacylglycerols) is the most valuable product of the seeds or fruits and other compounds such as starch, protein and fiber is  
 40 regarded as by-products with less value. Enhancing the quantity of oil per weight basis at the expense of other compounds in oil crops would therefore increase the value of the crop. If proteins that promote the allocation of reduced carbon into the production of oil can be up regulated by overexpression, the cells will  
 45 accumulate more oil at the expense of other products. This approach could not only be used to increase the oil content in already high oil producing organisms such as oil crops, they



could also lead to significant oil production in moderate or low oil containing crops such as soy, oat, maize, potato, sugar beats, and turnips as well as in microorganisms.

- 5 Increasing the oil content in plants and, in particular, in plant seeds is of great interest for traditional and modern plant breeding and in particular for plant biotechnology. Owing to the increasing consumption of vegetable oils for nutrition or industrial applications, possibilities of increasing or modifying vegetable oils are increasingly the subject of current research (for example Töpfer et al. (1995) Science 268:681-686). Its aim is in particular increasing the fatty acid content in seed oils.

- The fatty acids which can be obtained from the vegetable oils are also of particular interest. They are employed, for example, as bases for plasticizers, lubricants, surfactants, cosmetics and the like and are employed as valuable bases in the food and feed industries. Thus, for example, it is of particular interest to provide rapeseed oils with fatty acids with medium chain length since these are in demand in particular in the production of surfactants.

- The targeted modulation of plant metabolic pathways by recombinant methods allows the modification of the plant metabolism in an advantageous manner which, when using traditional breeding methods, could only be achieved after a complicated procedure or not at all. Thus, unusual fatty acids, for example specific polyunsaturated fatty acids, are only synthesized in certain plants or not at all in plants and can therefore only be produced by expressing the relevant gene in transgenic plants (for example Millar et al. (2000) Trends Plant Sci 5:95-101).

- Triacylglycerides and other lipids are synthesized from fatty acids. Fatty acid biosynthesis and triacylglyceride biosynthesis can be considered as separate biosynthetic pathways owing to the compartmentalization, but as a single biosynthetic pathway in view of the end product. Lipid synthesis can be divided into two part-mechanisms, one which might be termed "prokaryotic" and another which may be termed "eukaryotic" (Browse et al. (1986) Biochemical J 235:25-31; Ohlrogge & Browse (1995) Plant Cell 7:957-970). The prokaryotic mechanism is localized in the plastids and encompasses the biosynthesis of the free fatty acids which are exported into the cytosol, where they enter the eukaryotic mechanism in the form of fatty acid acyl-CoA esters and are esterified with glycerol-3-phosphate (G3P) to give phosphatidic acid (PA). PA is the starting point for the synthesis of neutral and polar lipids. The neutral lipids are



3

synthesized on the endoplasmic reticulum via the Kennedy pathway (Voelker (1996) Genetic Engineering, Setlow (ed.) 18:111-113; Shankline & Cahoon (1998) Annu Rev Plant Physiol Plant Mol Biol 49:611-649; Frentzen (1998) Lipids 100:161-166).

5

The last step in the synthesis of triacylglycerols has been shown to occur by two different enzymatic reactions, an acyl-CoA dependent reaction catalyzed by an acyl-CoA : diacylglycerol acyltransferase (Cases, et al., 1998; Lardizabal, et al., 2001) and the acyl-CoA independent reaction catalyzed by an phospholipid : diacylglycerol acyltransferase (Dahlgvist, et al., 2000). Two unrelated gene families encoding acyl-CoA : diacylglycerol acyltransferases have been identified in plants, animals and yeast, whereas the gene family encoding the acyl-CoA independent enzyme has been identified in yeast but not in plants or animals. In yeast, a total of four genes (are1, are2, lro1, dgal) belong to these three gene families, and they are the only genes known to contribute directly to triacylglycerol synthesis. Thus, no synthesis of triacylglycerol could be detected in yeast cells where all four genes were disrupted. In the present invention we show, that a fifth gene is present in yeast, which enhances the amount of triacylglycerol that accumulates in wildtype yeast.

It is an object of the present invention to provide alternative methods for increasing the oil content in plants.

We have found that this object is achieved by the present invention.

- A first subject matter of the invention comprises a method of increasing the total oil content in a plant organism or a tissue, organ, part, cell or propagation material thereof, comprising
- the transgenic expression of yeast TEP in said plant organism or in a tissue, organ, part, cell or propagation material thereof, and
  - the selection of plant organisms in which - in contrast to or comparison with the starting organism - the total oil content in said plant organism or in a tissue, organ, part, cell or propagation material thereof is increased.

Other proteins resulting in the same effect as the protein set forth in SEQ ID NO. 2 are obtainable from the specific sequences provided herein. Furthermore, it will be apparent that one can obtain natural and synthetic TEPs, including those with modified amino acid sequences and starting materials for synthetic-protein



4

modeling from the exemplified TEPs and from TEPs which are obtained through the use of such exemplified sequences. Modified amino acid sequences include sequences that have been mutated, truncated, increased and the like, whether such sequences were partially or wholly synthesized.

Further, the nucleic acid probes (DNA or RNA) derived from the SEQ-ID No. 1 of the present invention can be used to screen and recover "homologous" or "related" sequences from a variety of plant and microbial sources.

The present invention can be essentially characterized by the following aspects:

Example 1 shows the reduction of triacylglycerol accumulation in yeast cells lacking the YJR098c gene.

Example 2 shows the increased accumulation of triacylglycerol in yeast cells expressing the YJR098c gene in combination with a strong promoter.

Example 3 shows a significantly higher total oil content in the seeds of transgenic plant lines with increased expression of the YJR098c gene construct.

Use of a nucleic acid sequence SEQ-ID No: 1, encoding a protein SEQ-ID No: 2 that enhances the production of triacylglycerol (TAG), by genetic transformation of an oil-producing organism with said sequence in order to be expressed in this organism, resulting in an active protein that increases the oil content of the organism. The nucleic acid sequence is derived from the sequence shown in SEQ ID NO. 1 from the *Saccharomyces cerevisiae* YJR098c gene (genomic clone or cDNA) or from a nucleic acid sequence or cDNA that contains a nucleotide sequence coding for a protein with an amino acid sequence that is 60% or more identical to the amino acid sequence as presented in SEQ ID No: 2.

The gene product, which we refer to as a TAG synthesis enhancing protein (TEP) is most likely not itself catalyzing the synthesis of TAG, but its presence elevates the amount of TAG synthesized by other enzymes.

The instant invention pertains to a gene construct comprising a said nucleotide sequence SEQ ID No: 1 of the instant invention, which is operably linked to a heterologous nucleic acid.



5

The term operably linked means a serial organization e.g. of a promoter, coding sequence, terminator and/or further regulatory elements whereby each element can fulfill its original function during expression of the nucleotide sequence.

5

Further, a vector comprising the said nucleotide sequence SEQ ID No: 1 of the instant invention is contemplated in the instant invention. This includes also an expression vector which can harbor a selectable marker gene and/or nucleotide sequences 10 for the replication in a host cell and/or the integration into the genome of the host cell.

Furthermore, this invention relates to a method for producing a TEP in a host cell or progeny thereof including genetically 15 engineered oil seeds, yeast and moulds or any other oil-accumulating organism, via the expression of a construct in the cell. Of particular interest is the expression of the nucleotide sequences of the present invention from transcription initiation regions that are preferentially expressed in plant 20 seed tissues. It is further contemplated that an artificial gene sequence encoding TEP may be synthesized, especially to provide plant-preferred codons. Cells containing a TEP as a result of the production of a TEP encoding sequence are also contemplated within the scope of the invention.

25

Further, the invention pertains a transgenic cell or organism containing a said nucleotide sequence and/or a said gene construct and/or a said vector. The object of the instant invention is further a transgenic cell or organism which is an 30 eucaryotic cell or organism. Preferably, the transgenic cell or organism is a yeast cell or a plant cell or a plant. The instant invention further pertains said transgenic cell or organism having an increased biosynthetic pathway for the production of substrates for the synthesis of triacylglycerol. A transgenic 35 cell or organism having increased oil content is also contemplated within the scope of this invention.

Further, the invention pertains a transgenic cell or organism wherein the activity of TEP is increased in said cell or 40 organism. The increased activity of TEP is characterized by an alteration in gene expression, catalytic activity and/or regulation of activity of the enzyme. Moreover, a transgenic cell or organism is included in the instant invention, wherein the increased biosynthetic pathway for the production of sub- 45 strates for the production of triacylglycerol is characterized



6

e.g. by the prevention of accumulation of undesirable fatty acids in the membrane lipids.

In a different embodiment, this invention also relates to methods of using a DNA sequence coding for a TEP for increasing the oil-content within the cells of different organisms.

Further, the invention makes possible a process for elevating the production of triacylglycerol, which comprises growing transgenic cells or organisms under conditions whereby the nucleotide sequence SEQ-ID No: 1 is expressed in order to produce an protein in these cells with the ability of enhancing the production of triacylglycerol.

Corresponding genes coding for TEP can be isolated from other organisms, especially yeast-type organisms, like e.g.

*Schizosaccharomyces pombe*, *Yarrowia lipolytica*, *Zygosaccharomyces rouxii*, *Saccharomyces cerevisiae*, *Emericella nidulans* and *Debaryomyces hansenii*.

Transgenic organisms comprising, in their genome or on a plasmid, a nucleic acid sequence SEQ ID No:1 according to the above, transferred by recombinant DNA technology. One important type of transgenic organism covered by this invention are commercially relevant plants in which said nucleotide sequence preferably would be expressed under the control of a storage organ specific promoter. Alternatively, the nucleotide sequence could also be expressed under the control of a seed-specific promoter or any other promoter suitable for tissue-specific high-level expression in plants.

A protein encoded by a DNA molecule according to SEQ ID NO. 1 or a functional biologically active fragment thereof having TEP activity in transgenic organisms. Alternatively, the protein produced in an organism, which has the amino acid sequence set forth in SEQ ID NO. 2 or an amino acid sequence with at least 60 % homology to said amino acid sequence having TEP activity. Preferably the protein is isolated from *Saccharomyces cerevisiae*.

Use of a protein according to SEQ ID No: 2 or derivatives of that protein having TEP activity for the increased production of triacylglycerols.

Surprisingly, it has been found that the heterologous expression of the yeast TEP from *Saccharomyces cerevisiae* SEQ ID NO: 1 in *Arabidopsis* leads to a significantly increased triacylglyceride



7

(storage oils) content in the seeds. The oil content was increased by approximately 5%, in one transgenic line even by 10%, compared with wild-type control plants. The transgenic expression of the yeast TEP had no adverse effects on the growth or other properties of the transformed plants.

The method according to the invention can be applied in principle to all plant species, in addition to the species *Arabidopsis thaliana*, which is employed as model plant. The method according to the invention is preferably applied to oil crops whose oil content is already naturally high and/or for the industrial production of oils.

"Plant" organism or tissue, organ, part, cell or propagation material thereof" is generally understood as meaning any single- or multi-celled organism or a cell, tissue, part or propagation material (such as seeds or fruit) of same which is capable of photosynthesis. Included for the purpose of the invention are all genera and species of higher and lower plants of the Plant Kingdom. Annual, perennial, monocotyledonous and dicotyledonous plants are preferred. Also included are mature plants, seeds, shoots and seedlings, and parts, propagation material (for example tubers, seeds or fruits) and cultures derived from them, for example cell cultures or callus cultures.

"Plant" encompasses all annual and perennial monocotyledonous or dicotyledonous plants and includes by way of example, but not by limitation, those of the genera Cucurbita, Rosa, Vitis, Juglans, Fragaria, Lotus, Medicago, Onobrychis, Trifolium, Trigonella, Vigna, Citrus, Linum, Geranium, Manihot, Daucus, Arabidopsis, Brassica, Raphanus, Sinapis, Atropa, Capsicum, Datura, Hyoscyamus, Lycopersicon, Nicotiana, Solarium, Petunia, Digitalis, Majorana, Cichorium, Helianthus, Lactuca, Bromus, Asparagus, Antirrhinum, Heterocallis, Nemesis, Pelargonium, Panieum, Pennisetum, Ranunculus, Senecio, Salpiglossis, Cucumis, Bro-waalia, Glycine, Pisum, Phaseolus, Lolium, Oryza, Zea, Avena, Hordeum, Secale, Triticum, Sorghum, Picea and Populus.

Preferred plants are those from the following plant families: Amaranthaceae, Asteraceae, Brassicaceae, Carophyllaceae, Chenopodiaceae, Compositae, Cruciferae, Cucurbitaceae, Labiatae, Leguminosae, Papilionoideae, Liliaceae, Linaceae, Malvaceae, Rosaceae, Rubiaceae, Saxifragaceae, Scrophulariaceae, Solanaceae, Sterculiaceae, Tetragoniaceae, Theaceae, Umbelliferae.

45



## 8

Preferred monocotyledonous plants are selected in particular from the monocotyledonous crop plants such as, for example, the Gramineae family, such as rice, maize, wheat or other cereal species such as barley, millet and sorghum, rye, triticales or 5 oats, and sugar cane, and all grass species.

The invention is applied very particularly preferably to dicotyledonous plant organisms. Preferred dicotyledonous plants are selected in particular from the dicotyledonous 10 crop plants such as, for example,

- Asteraceae such as *Helianthus annuus* (sunflower), *tagetes* or *calendula* and others,
- 15 - Compositae, especially the genus *Lactuca*, very particularly the species *sativa* (lettuce) and others,
- Cruciferae, particularly the genus *Brassica*, very particularly the species *napus* (oilseed rape), *campestris* (beet), 20 *oleracea* cv *Tastie* (cabbage), *oleracea* cv *Snowball Y* (cauliflower) and *oleracea* cv *Emperor* (broccoli) and other cabbages; and the genus *Arabidopsis*, very particularly the species *thaliana*, and cress or canola and others,
- 25 - Cucurbitaceae such as melon, pumpkin/squash or zucchini and others,
- Leguminosae, particularly the genus *Glycine*, very particularly the species *max* (soybean), *soya*, and *alfalfa*, 30 *pea*, *beans* or *peanut* and others,
- Rubiaceae, preferably the subclass *Lamiidae* such as, for example *Coffea arabica* or *Coffea liberica* (coffee bush) and others,
- 35 - Solanaceae, particularly the genus *Lycopersicon*, very particularly the species *esculentum* (tomato), the genus *Solanum*, very particularly the species *tuberosum* (potato) and *melongena* (aubergine) and the genus *Capsicum*, very 40 particularly the genus *annuum* (pepper) and tobacco or paprika and others,
- Sterculiaceae, preferably the subclass *Dilleniidae* such as, for example, *Theobroma cacao* (cacao bush) and others,

45



9

- Theaceae, preferably the subclass Dilleniidae such as, for example, *Camellia sinensis* or *Thea sinensis* (tea shrub) and others,
- 5 - Umbelliferae, particularly the genus *Daucus* (very particularly the species *carota* (carrot)) and *Apium* (very particularly the species *graveolens dulce* (celery)) and others;
- 10 and linseed, cotton, hemp, flax, cucumber, spinach, carrot, sugar beet and the various tree, nut and grapevine species, in particular banana and kiwi fruit.

Also encompassed are ornamental plants, useful or ornamental  
 15 trees, flowers, cut flowers, shrubs or turf plants which may be mentioned by way of example but not by limitation are angiosperms, bryophytes such as, for example, Hepaticae (liverworts) and Musci (mosses); pteridophytes such as ferns, horsetail and clubmosses; gymnosperms such as conifers, cycades, ginkgo  
 20 and Gnetatae; algae such as Chlorophyceae, Phaeophyceae, Rhodophyceae, Myxophyceae, Xanthophyceae, Bacillariophyceae (diatoms) and Euglenophyceae. Plants within the scope of the invention comprise by way of example and not by way of limitation, the families of the Rosaceae such as rose, Ericaceae such as rhododendron and azalea, Euphorbiaceae such as poinsettias and croton,  
 25 Caryophyllaceae such as pinks, Solanaceae such as petunias, Gesneriaceae such as African violet, Balsaminaceae such as touch-me-not, Orchidaceae such as orchids, Iridaceae such as gladioli, iris, freesia and crocus, Compositae such as marigold, Gerania-  
 30 ceae such as geranium, Liliaceae such as dracena, Moraceae such as ficus, Araceae such as cheeseplant and many others.

Furthermore, plant organisms for the purposes of the invention are further organisms capable of being photosynthetically active  
 35 such as, for example, algae, cyanobacteria and mosses. Preferred algae are green algae such as, for example, algae from the genus *Haematococcus*, *Phaedactylum tricornutum*, *Volvox* or *Dunaliella*. *Synechocystis* is particularly preferred.

40 Most preferred are oil crops. Oil crops are understood as being plants whose oil content is already naturally high and/or which can be used for the industrial production of oils. These plants can have a high oil content and/or else a particular fatty acid composition which is of interest industrially. Preferred plants  
 45 are those with a lipid content of at least 1% by weight. Oil crops encompassed by way of example: *Borvago officinalis* (borage); *Brassica* species such as *B. campestris*, *B. napus*,



## 10

*B. rapa* (mustard, oilseed rape or turnip rape); *Cannabis sativa* (hemp); *Carthamus tinctorius* (safflower); *Cocos nucifera* (coconut); *Crambe abyssinica* (crambe); *Cuphea* species (*Cuphea* species yield fatty acids of medium chain length, in particular for industrial applications); *Elaeis guineensis* (African oil palm); *Elaeis oleifera* (American oil palm); *Glycine max* (soybean); *Gossypium hirsutum* (American cotton); *Gossypium barbadense* (Egyptian cotton); *Gossypium herbaceum* (Asian cotton); *Helianthus annuus* (sunflower); *Linum usitatissimum* (linseed or flax); *Oenothera biennis* (evening primrose); *Olea europaea* (olive); *Oryza sativa* (rice); *Ricinus communis* (castor); *Sesamum indicum* (sesame); *Triticum* species (wheat); *Zea mays* (maize), and various nut species such as, for example, walnut or almond.

15 "Total oil content" refers to the sum of all oils, preferably to the sum of the triacylglycerides.

"Oils" encompasses neutral and/or polar lipids and mixtures of these. Those mentioned in Table 1 may be mentioned by way of example, but not by limitation.

Table 1: Classes of plant lipids

25	Neutrals lipids	Triacylglycerol (TAG)
		Diacylglycerol (DAG)
		Monoacylglycerol (MAG)
30	Polar lipids	Monogalactosyldiacylglycerol (MGDG)
		Digalactosyldiacylglycerol (DGDG)
		Phosphatidylglycerol (PG)
		Phosphatidylcholine (PC)
		Phosphatidylethanolamine (PE)
		Phosphatidylinositol (PI)
		Phosphatidylserine (PS)
35		Sulfoquinovosyldiacylglycerol

Neutral lipids preferably refers to triacylglycerides. Both neutral and polar lipids may comprise a wide range of various fatty acids. The fatty acids mentioned in Table 2 may be mentioned by way of example, but not by limitation.



## 11

Table 2: Overview over various fatty acids (selection)

<sup>1</sup> Chain length: number of double bonds

\* not naturally occurring in plants

	Nomenclature <sup>1</sup>	Name
5	16:0	Palmitic acid
	16:1	Palmitoleic acid
	16:3	Roughanic acid
10	18:0	Stearic acid
	18:1	Oleic acid
	18:2	Linoleic acid
	18:3	Linolenic acid
	$\gamma$ -18:3-18:3	Gamma-linolenic acid *
15	20:0	Arachidic acid
	22:6	Docosahexaenoic acid (DHA) *
	20:2	Eicosadienoic acid
	20:4	Arachidonic acid (AA) *
	20:5	Eicosapentaenoic acid (EPA) *
	22:1	Erucic acid

20

Oils preferably relates to seed oils.

"Increase in" the total oil content refers to the increased oil content in a plant or a part, tissue or organ thereof,

25 preferably in the seed organs of the plants. In this context, the oil content is at least 5%, preferably at least 10%, particularly preferably at least 15%, very particularly preferably at least 20%, most preferably at least 25% increased under otherwise

identical conditions in comparison with a starting plant which

30 has not been subjected to the method according to the invention, but is otherwise unmodified. Conditions in this context means all of the conditions which are relevant for germination, culture or growth of the plant, such as soil conditions, climatic conditions, light conditions, fertilization, irrigation, plant  
35 protection treatment and the like.

"Yeast TEP" generally refers to all those proteins which are capable of increasing the oil content in oil producing organisms, especially microorganisms, yeast, fungi and plants and are  
40 identical to SEQ ID No: 2 or have homology to SEQ ID No: 2.

Yeast refers to the group of unicellular fungi with a pronounced cell wall and formation of pseudomycelium (in contrast to molds). They reproduce vegetatively by budding and/or fission (Schizo-  
45 saccharomyces and Saccharomycodes, respectively).



## 12

Encompassed are what are known as false yeasts, preferably the families Cryptococcaceae, Sporobolomycetaceae with the genera Cryptococcus, Torulopsis, Pityrosporum, Brettanomyces, Candida, Klöckera, Trigonopsis, Trichosporon, Rhodotorula and Sporobolomyces and Bullera, and true yeasts (yeasts which also reproduce sexually; ascus), preferably the families endo- and saccharomycetaceae, with the genera Saccharomyces, Debaromyces, Lipomyces, Hansenula, Endomycopsis, Pichia, Hanseniaspora. Most preferred are the genera Saccharomyces cerevisiae, Pichia pastoris, Hansenula polymorpha, Schizosaccharomyces pombe, Kluyveromyces lactis, Zygosaccharomyces rouxii, und Yarrowia lipolitica, Emericella nidulans, Aspergillus nidulans, Debaryomyces hansenii and Torulaspora hansenii.

15 Yeast TEP refers in particular to the polypeptide sequence SEQ ID No. 2.

Most preferably, yeast TEP refers to the yeast protein TEP as shown in SEQ ID NO: 2 and functional equivalents or else  
20 functionally equivalent portions of the above.

Functional equivalents refers in particular to natural or artificial mutations of the yeast protein TEP as shown in SEQ ID NO: 2 and homologous polypeptides from other yeasts which  
25 have the same essential characteristics of a yeast TEP as defined above. Mutations encompass substitutions, additions, deletions, inversions or insertions of one or more amino acid residues.

The yeast TEP to be employed advantageously within the scope of  
30 the present invention can be found readily by database searches or by screening gene or cDNA libraries using the yeast TEP sequence shown in SEQ ID NO: 2, which is given by way of example, or the nucleic acid sequence as shown in SEQ ID NO: 1, which encodes the latter, as search sequence or probe.

35 Said functional equivalents preferably have at least 60%, particularly preferably at least 70%, particularly preferably at least 80%, most preferably at least 90% homology with the protein of SEQ ID NO: 2.

40 Homology between two polypeptides is understood as meaning the identity of the amino acid sequence over the entire sequence length which is calculated by comparison with the aid of the program algorithm GAP (Wisconsin Package Version 10.0, University  
45 of Wisconsin, Genetics Computer Group (GCG), Madison, USA), setting the following parameters:



13

Gap Weight: 8

Length Weight: 2

Average Match: 2,912

Average Mismatch: -2,003

5 For example, a sequence with at least 80% homology with the sequence SEQ ID NO: 2 at the protein level is understood as meaning a sequence which, upon comparison with the sequence SEQ ID NO: 2 with the above program algorithm and the above parameter set has at least 80% homology.

10

Functional equivalents also encompass those proteins which are encoded by nucleic acid sequences which have at least 60%, particularly preferably at least 70%, particularly preferably at least 80%, most preferably at least 90% homology with the

15 nucleic acid sequence with the SEQ ID NO: 1.

Homology between two nucleic acid sequences is understood as meaning the identity of the two nucleic acid sequences over the entire sequence length which is calculated by comparison with the

20 aid of the program algorithm GAP (Wisconsin Package Version 10.0, University of Wisconsin, Genetics Computer Group (GCG), Madison, USA), setting the following parameters:

Gap Weight: 50

Length Weight: 3

25

Average Match: 10

Average Mismatch: 0

For example, a sequence which has at least 80% homology with the sequence SEQ ID NO: 1 at the nucleic acid level is understood

30 as meaning a sequence which, upon comparison with the sequence SEQ ID NO: 1 within the above program algorithm with the above parameter set has a homology of at least 80%.

Functional equivalents also encompass those proteins which  
35 are encoded by nucleic acid sequences which hybridize under standard conditions with a nucleic acid sequence described by SEQ ID NO: 1, the nucleic acid sequence which is complementary thereto or parts of the above and which have the essential characteristics for a yeast TEP.

40

"Standard hybridization conditions" is to be understood in the broad sense, but preferably refers to stringent hybridization conditions. Such hybridization conditions are described, for example, by Sambrook J, Fritsch EF, Maniatis T et al., in Molecular Cloning (A Laboratory Manual), 2nd edition, Cold Spring Harbor Laboratory Press, 1989, pages 9.31-9.57) or in Current Protocols in Molecular Biology, John Wiley & Sons, N.Y. (1989),

45



14

6.3.1-6.3.6. For example, the conditions during the wash step can be selected from the range of high-stringency conditions (with approximately 0.2X SSC at 50°C, preferably at 65°C) (20X SSC: 0.3 M sodium citrate, 3 M NaCl, pH 7.0). Denaturing agents such as, for example, formamide or SDS may also be employed during hybridization. In the presence of 50% formamide, hybridization is preferably carried out at 42°C.

The invention furthermore relates to transgenic expression constructs which can ensure a transgenic expression of a yeast TEP in a plant organism or a tissue, organ, part, cells or propagation material of said plant organism.

The definition given above applies to yeast TEP, with the transgenic expression of a yeast TEP described by the sequence with the SEQ ID NO: 2 being particularly preferred.

In said transgenic expression constructs, a nucleic acid molecule encoding a yeast TEP is preferably in operable linkage with at least one genetic control element (for example a promoter) which ensures expression in a plant organism or a tissue, organ, part, cell or propagation material of same.

Especially preferred are transgenic expression cassettes wherein the nucleic acid sequence encoding a TEP is described by

- a) a sequence with the SEQ ID NO: 1,
- b) a sequence derived from a sequence with the SEQ ID NO: 1 in accordance with the degeneracy of the genetic code
- c) a sequence which has at least 60% identity with the sequence with the SEQ ID NO: 1.

Operable linkage is understood as meaning, for example, the sequential arrangement of a promoter with the nucleic acid sequence encoding a yeast TEP which is to be expressed (for example the sequence as shown in SEQ ID NO: 1 and, if appropriate, further regulatory elements such as, for example, a terminator in such a way that each of the regulatory elements can fulfil its function when the nucleic acid sequence is expressed recombinantly. Direct linkage in the chemical sense is not necessarily required for this purpose. Genetic control sequences such as, for example, enhancer sequences can also exert their function on the target sequence from positions which are further removed or indeed from other DNA molecules. Preferred arrangements are those in which the nucleic acid sequence to be



15

expressed recombinantly is positioned behind the sequence acting as promoter so that the two sequences are linked covalently to each other. The distance between the promoter sequence and the nucleic acid sequence to be expressed recombinantly is preferably less than 200 base pairs, particularly preferably less than 100 base pairs, very particularly preferably less than 50 base pairs.

Operable linkage and a transgenic expression cassette can both be effected by means of conventional recombination and cloning techniques as they are described, for example, in Maniatis T, Fritsch EF and Sambrook J (1989) Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor (NY), in Silhavy TJ, Berman ML und Enquist LW (1984) Experiments with Gene Fusions, Cold Spring Harbor Laboratory, Cold Spring Harbor (NY), in Ausubel FM et al. (1987) Current Protocols in Molecular Biology, Greene Publishing Assoc. and Wiley Interscience and in Gelvin et al. (1990) In: Plant Molecular Biology Manual. However, further sequences which, for example, act as a linker with specific cleavage sites for restriction enzymes, or of a signal peptide, may also be positioned between the two sequences. Also, the insertion of sequences may lead to the expression of fusion proteins. Preferably, the expression cassette composed of a promoter linked to a nucleic acid sequence to be expressed can be in a vector-integrated form and can be inserted into a plant genome, for example by transformation.

However, a transgenic expression cassette is also understood as meaning those constructs where the nucleic acid sequence encoding a yeast TEP is placed behind an endogenous plant promoter in such a way that the latter brings about the expression of the yeast TEP.

Promoters which are preferably introduced into the transgenic expression cassettes are those which are operable in a plant organism or a tissue, organ, part, cell or propagation material of same. Promoters which are operable in plant organisms is understood as meaning any promoter which is capable of governing the expression of genes, in particular foreign genes, in plants or plant parts, plant cells, plant tissues or plant cultures. In this context, expression may be, for example, constitutive, inducible or development-dependent.

45



The following are preferred:

a) Constitutive promoters

5 "Constitutive" promoters refers to those promoters which ensure expression in a large number of, preferably all, tissues over a substantial period of plant development, preferably at all times during plant development (Benfey et al. (1989) EMBO J 8:2195-2202). A plant promoter or  
10 promoter originating from a plant virus is especially preferably used. The promoter of the CaMV (cauliflower mosaic virus) 35S transcript (Franck et al. (1980) Cell 21:285-294; Odell et al. (1985) Nature 313:810-812; Shewmaker et al. (1985) Virology 140:281-288; Gardner et al. (1986) Plant  
15 Mol Biol 6:221-228) or the 19S CaMV promoter (US 5,352,605; WO 84/02913; Benfey et al. (1989) EMBO J 8:2195-2202) are especially preferred. Another suitable constitutive promoter is the Rubisco small subunit (SSU) promoter (US 4,962,028), the leguminB promoter (GenBank Acc. No. X03677), the promoter  
20 of the nopaline synthase from Agrobacterium, the TR dual promoter, the OCS (octopine synthase) promoter from Agrobacterium, the ubiquitin promoter (Holtorf S et al. (1995) Plant Mol Biol 29:637-649), the ubiquitin 1 promoter (Christensen et al. (1992) Plant Mol Biol 18:675-689; Bruce et al. (1989) Proc Natl Acad Sci USA 86:9692-9696), the Smas promoter;  
25 the cinnamyl alcohol dehydrogenase promoter (US 5,683,439), the promoters of the vacuolar ATPase subunits, the promoter of the Arabidopsis thaliana nitrilase-1 gene (GenBank Acc. No. U38846, nucleotides 3862 to 5325 or else 5342) or the  
30 promoter of a proline-rich protein from wheat (WO 91/13991), and further promoters of genes whose constitutive expression in plants is known to the skilled worker. The CaMV 35S promoter and the Arabidopsis thaliana nitrilase-1 promoter are particularly preferred.

35

b) Tissue-specific promoters

Furthermore preferred are promoters with specificities for seeds, such as, for example, the phaseolin promoter  
40 (US 5,504,200; Bustos MM et al. (1989) Plant Cell 1(9):839-53), the promoter of the 2S albumin gene (Joseffson LG et al. (1987) J Biol Chem 262:12196-12201), the legumine promoter (Shirsat A et al. (1989) Mol Gen Genet 215(2):326-331), the USP (unknown seed protein) promoter  
45 (Bäumlein H et al. (1991) Mol Gen Genet 225(3):459-67), the napin gene promoter (US 5,608,152; Stalberg K et al. (1996) L Planta 199:515-519), the promoter of the sucrose binding



17

proteins (WO 00/26388) or the legumin B4 promoter (LeB4; Bäumlein H et al. (1991) Mol Gen Genet 225: 121-128; Bäumlein et al. (1992) Plant Journal 2(2):233-9; Fiedler U et al. (1995) Biotechnology (NY) 13(10):1090f), the Arabidopsis oleosin promoter (WO 98/45461), and the Brassica Bce4 promoter (WO 91/13980).

Further suitable seed-specific promoters are those of the gene encoding high-molecular weight glutenin (HMWG), gliadin, branching enzyme, ADP glucose pyrophosphatase (AGPase) or starch synthase. Promoters which are furthermore preferred are those which permit a seed-specific expression in monocots such as maize, barley, wheat, rye, rice and the like. The promoter of the lpt2 or lpt1 gene (WO 95/15389, WO 95/23230) or the promoters described in WO 99/16890 (promoters of the hordein gene, the glutelin gene, the oryzin gene, the prolamins gene, the gliadin gene, the glutelin gene, the zein gene, the casirin gene or the secalin gene) can advantageously be employed.

c) Chemically inducible promoters

The expression cassettes may also contain a chemically inducible promoter (review article: Gatz et al. (1997) Annu Rev Plant Physiol Plant Mol Biol 48:89-108), by means of which the expression of the exogenous gene in the plant can be controlled at a particular point in time. Such promoters such as, for example, the PRP1 promoter (Ward et al. (1993) Plant Mol Biol 22:361-366), a salicylic acid-inducible promoter (WO 95/19443), a benzenesulfonamide-inducible promoter (EP 0 388 186), a tetracyclin-inducible promoter (Gatz et al. (1992) Plant J 2:397-404), an abscisic acid-inducible promoter (EP 0 335 528) or an ethanol-cyclohexanone-inducible promoter (WO 93/21334) can likewise be used. Also suitable is the promoter of the glutathione-S transferase isoform II gene (GST-II-27), which can be activated by exogenously applied safeners such as, for example, N,N-diallyl-2,2-dichloroacetamide (WO 93/01294) and which is operable in a large number of tissues of both monocots and dicots.

Particularly preferred are constitutive promoters, very particularly preferred seed-specific promoters, in particular the napin promoter and the USP promoter.



In addition, further promoters which make possible expression in further plant tissues or in other organisms such as, for example, E.coli bacteria, may be linked operably with the nucleic acid sequence to be expressed. Suitable plant promoters are, in principle, all of the above-described promoters.

The nucleic acid sequences present in the transgenic expression cassettes according to the invention or transgenic vectors can be linked operably with further genetic control sequences besides a promoter. The term genetic control sequences is to be understood in the broad sense and refers to all those sequences which have an effect on the establishment or the function of the expression cassette according to the invention. Genetic control sequences modify, for example, transcription and translation in prokaryotic or eukaryotic organisms. The transgenic expression cassettes according to the invention preferably encompass a plant-specific promoter 5'-upstream of the nucleic acid sequence to be expressed recombinantly in each case and, as additional genetic control sequence, a terminator sequence 3'-downstream, and, if appropriate, further customary regulatory elements, in each case linked operably with the nucleic acid sequence to be expressed recombinantly.

Genetic control sequences also encompass further promoters, promoter elements or minimal promoters capable of modifying the expression-controlling properties. Thus, genetic control sequences can, for example, bring about tissue-specific expression which is additionally dependent on certain stress factors. Such elements are, for example, described for water stress, abscisic acid (Lam E and Chua NH, J Biol Chem 1991; 266(26): 17131-17135) and thermal stress (Schoffl F et al. (1989) Mol Gen Genetics 217(2-3):246-53).

Further advantageous control sequences are, for example, in the Gram-positive promoters amy and SPO2, and in the yeast or fungal promoters ADC1, MFA, AC, P-60, CYC1, GAPDH, TEF, rp28, ADH.

In principle all natural promoters with their regulatory sequences like those mentioned above may be used for the method according to the invention. In addition, synthetic promoters may also be used advantageously.

Genetic control sequences further also encompass the 5'-untranslated regions, introns or nonencoding 3'-region of genes, such as, for example, the actin-1 intron, or the Adh1-S intron 1, 2 and 6 (for general reference, see: The Maize Handbook, Chapter 116, Freeling and Walbot, Eds., Springer, New York (1994)). It



19

has been demonstrated that these may play a significant role in regulating gene expression. Thus, it has been demonstrated that 5'-untranslated sequences can enhance the transient expression of heterologous genes. Translation enhancers which may be mentioned by way of example are the tobacco mosaic virus 5' leader sequence (Gallie et al. (1987) Nucl Acids Res 15:8693-8711) and the like. They may furthermore promote tissue specificity (Rouster J et al. (1998) Plant J 15:435-440).

- 10 The transient expression cassette can advantageously contain one or more of what are known as enhancer sequences in operable linkage with the promoter, and these make possible an increased recombinant expression of the nucleic acid sequence. Additional advantageous sequences such as further regulatory elements or
- 15 terminators may also be inserted at the 3' end of the nucleic acid sequences to be expressed recombinantly. One or more copies of the nucleic acid sequences to be expressed recombinantly may be present in the gene construct.
- 20 Polyadenylation signals which are suitable as control sequences are plant polyadenylation signals, preferably those which correspond essentially to Agrobacterium tumefaciens T-DNA polyadenylation signals, in particular those of gene 3 of the T-DNA (octopine synthase) of the Ti plasmid pTiACH5 (Gielen et al. (1984) EMBO J 3:835 et seq.) or functional equivalents thereof. ....
- 25 Examples of particularly suitable terminator sequences are the OCS (octopine synthase) terminator and the NOS (nopaline synthase) terminator.
- 30 Control sequences are furthermore understood as those which make possible homologous recombination or insertion into the genome of a host organism, or removal from the genome. In the case of homologous recombination, for example, the coding sequence of the specific endogenous gene can be exchanged in a directed fashion
- 35 for a sequence encoding a dsRNA. Methods such as the cre/lox technology permit the tissue-specific, possibly inducible, removal of the expression cassette from the genome of the host organism (Sauer B (1998) Methods. 14(4):381-92). Here, certain flanking sequences are added to the target gene (lox sequences),
- 40 and these make possible removal by means of cre recombinase at a later point in time.

A recombinant expression cassette and the recombinant vectors derived from it may comprise further functional elements.

- 45 The term functional element is to be understood in the broad sense and refers to all those elements which have an effect on generation, replication or function of the expression cassettes,



vectors or transgenic organisms according to the invention. Examples which may be mentioned, but not by way of limitation, are:

- 5 a) Selection markers which confer resistance to a metabolism inhibitor such as 2-deoxyglucose-6-phosphate (WO 98/45456), antibiotics or biocides, preferably herbicides, such as, for example, kanamycin, G 418, bleomycin, hygromycin, or phosphothricin and the like. Particularly preferred  
10 selection markers are those which confer resistance to herbicides. The following may be mentioned by way of example: DNA sequences which encode phosphothricin acetyltransferases (PAT) and which inactivate glutamine synthase inhibitors (bar and pat gene), 5-enolpyruvylshikimate-3-phosphate synthase  
15 genes (EPSP synthase genes), which confer resistance to Glyphosate (N-(phosphonomethyl)glycine), the gox gene, which encodes Glyphosate-degrading enzyme (Glyphosate oxidoreductase), the deh gene (encoding a dehalogenase which inactivates dalapon), sulfonylurea- and imidazolinone-  
20 inactivating acetolactate synthases, and bxn genes which encode nitrilase enzymes which degrade bromoxynil, the aasa gene, which confers resistance to the antibiotic spectinomycin, the streptomycin phosphotransferase (SPT) gene, which permits resistance to streptomycin, the neomycin phosphotransferase (NPTII) gene, which confers resistance to kana-  
25 mycin or geneticidin, the hygromycin phosphotransferase (HPT) gene, which confers resistance to hygromycin, the acetolactate synthase gene (ALS), which confers resistance to sulfonylurea herbicides, (for example, mutated ALS variants with, for example, the S4 and/or Hra mutation).  
30
- b) Reporter genes which encode readily quantifiable proteins and which allow the transformation efficacy or the expression site or time to be assessed via their color or enzyme activity. Very particularly preferred in this context are reporter  
35 proteins (Schenborn E, Groskreutz D. Mol Biotechnol. 1999; 13(1):29-44) such as the "green fluorescent protein" (GFP) (Sheen et al. (1995) Plant Journal 8(5):777-784), chloramphenicol transferase, a luciferase (Ow et al. (1986) Science 234:856-859), the aequorin gene (Prasher et al. (1985) Biochem Biophys Res Commun 126(3):1259-1268),  $\beta$ -galactosidase, with  $\beta$ -glucuronidase being very particularly preferred (Jefferson et al. (1987) EMBO J 6:3901-3907).  
40
- 45 c) Replication origins which allow replication of the expression cassettes or vectors according to the invention in, for example, E.coli. Examples which may be mentioned are ORI



(origin of DNA replication), the pBR322 ori or the P15A ori (Sambrook et al.: Molecular Cloning. A Laboratory Manual, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989).

5

d) Elements which are required for agrobacterium-mediated plant transformation such as, for example, the right or left border of the T-DNA, or the vir region.

- 10 To select cells which have successfully undergone homologous recombination or else cells which have successfully been transformed, it is generally required additionally to introduce a selectable marker which confers resistance to a biocide (for example a herbicide), a metabolism inhibitor such as 2-deoxy-  
15 glucose-6-phosphate (WO 98/45456) or an antibiotic to the cells which have successfully undergone recombination. The selection marker permits the selection of the transformed cells from untransformed cells (McCormick et al. (1986) Plant Cell Reports 5:81-84).

20

In addition, said recombinant expression cassette or vectors may comprise further nucleic acid sequences which do not encode a yeast TEP and whose recombinant expression leads to a further increase in fatty acid biosynthesis. By way of example, but  
25 not by limitation, such a proOIL nucleic acid sequence which is additionally expressed recombinantly can be selected from among nucleic acids encoding acetyl-CoA carboxylase (ACCase), glycerol-3-phosphate acyltransferase (GPAT), lysophosphatidate acyltransferase (LPAT), diacylglycerol acyltransferase (DAGAT)  
30 and phospholipid:diacylglycerol acyltransferase (PDAT). Such sequences are known to the skilled worker and are readily accessible from databases or suitable cDNA libraries of the respective plants.

- 35 An expression cassette according to the invention can advantageously be introduced into an organism or cells, tissues, organs, parts or seeds thereof (preferably into plants or plant cells, tissues, organs, parts or seeds) by using vectors in which the recombinant expression cassettes are present. The invention  
40 therefore furthermore relates to said recombinant vectors which encompass a recombinant expression cassette for a yeast TEP.

For example, vectors may be plasmids, cosmids, phages, viruses or else agrobacteria. The expression cassette can be introduced  
45 into the vector (preferably a plasmid vector) via a suitable restriction cleavage site. The resulting vector is first introduced into E.coli. Correctly transformed E.coli are selected,



The invention furthermore relates to transgenic plant organisms or tissues, organs, parts, cells or propagation material thereof which comprise a yeast TEP as defined above, a transgenic expression cassette for a yeast TEP or a transgenic vector encompassing such an expression cassette.

Such a transgenic plant organism is generated, for example, by means of transformation or transfection of the corresponding proteins or nucleic acids. The generation of a transformed organism (or a transformed cell or tissue) requires introducing the DNA in question (for example the expression vector), RNA or protein into the host cell in question. A multiplicity of methods is available for this procedure, which is termed transformation (or transduction or transfection) (Keown et al. (1990) *Methods in Enzymology* 185:527-537). Thus, the DNA or RNA can be introduced for example directly by microinjection or by bombardment with DNA-coated microparticles. The cell may also be permeabilized chemically, for example with polyethylene glycol, so that the DNA may reach the cell by diffusion. The DNA can also be introduced by protoplast fusion with other DNA-comprising units such as minicells, cells, lysosomes or liposomes. Electroporation is a further suitable method for introducing DNA; here, the cells are permeabilized reversibly by an electrical pulse. Soaking plant parts in DNA solutions, and pollen or pollen tube transformation, are also possible. Such methods have been described (for example in Bilang et al. (1991) *Gene* 100:247-250; Scheid et al. (1991) *Mol Gen Genet* 228:104-112; Guerche et al. (1987) *Plant Science* 52:111-116; Neuhaus et al. (1987) *Theor Appl Genet* 75:30-36; Klein et al. (1987) *Nature* 327:70-73; Howell et al. (1980) *Science* 208:1265; Horsch et al. (1985) *Science* 227:1229-1231; DeBlock et al. (1989) *Plant Physiology* 91:694-701; *Methods for Plant Molecular Biology* (Weissbach and Weissbach, eds.) Academic Press Inc. (1988); and *Methods in Plant Molecular Biology* (Schuler and Zielinski, eds.) Academic Press Inc. (1989)).

In plants, the methods which have been described for transforming and regenerating plants from plant tissues or plant cells are exploited for transient or stable transformation. Suitable  
45 methods are, in particular, protoplast transformation by polyethylene glycol-induced DNA uptake, the biolistic method with the gene gun, what is known as the particle bombardment method.



electroporation, the incubation of dry embryos in DNA-containing solution, and microinjection.

In addition to these "direct" transformation techniques, transformation may also be effected by bacterial infection by means of *Agrobacterium tumefaciens* or *Agrobacterium rhizogenes* and the transfer of corresponding recombinant Ti plasmids or Ri plasmids by infection with transgenic plant viruses. *Agrobacterium*-mediated transformation is best suited to cells of dicotyledonous plants. The methods are described, for example, in Horsch RB et al. (1985) *Science* 225: 1229f).

When *agrobacteria* are used, the expression cassette is to be integrated into specific plasmids, either into a shuttle vector or into a binary vector. If a Ti or Ri plasmid is to be used for the transformation, at least the right border, but in most cases the right and left border, of the Ti or Ri plasmid T-DNA is linked to the expression cassette to be introduced as flanking region.

Binary vectors are preferably used. Binary vectors are capable of replication both in *E. coli* and in *Agrobacterium*. As a rule, they contain a selection marker gene and a linker or polylinker flanked by the right and left T-DNA border sequence. They can be transformed directly into *Agrobacterium* (Holsters et al. (1978) *Mol. Gen. Genet.* 163:181-187). The selection marker gene, which is, for example, the *nptII* gene, which confers resistance to kanamycin, permits a selection of transformed *agrobacteria*. The *Agrobacterium* which acts as host organism in this case should already contain a plasmid with the *vir* region. The latter is required for transferring the T-DNA to the plant cells. An *Agrobacterium* transformed in this way can be used for transforming plant cells. The use of T-DNA for the transformation of plant cells has been studied intensively and described (EP 120 516; Hoekema, In: The Binary Plant Vector System, Offsetdrukkerij Kanthers B.V., Alblasserdam, Chapter V; An et al. (1985) *EMBO J* 4:277-287). Various binary vectors, some of which are commercially available, such as, for example, pBI101.2 or pBIN19 (Clontech Laboratories, Inc. USA), are known.

Further promoters which are suitable for expression in plants have been described (Rogers et al. (1987) *Meth. in Enzymol.* 153:253-277; Schardl et al. (1987) *Gene* 61:1-11; Berger et al. (1989) *Proc. Natl. Acad. Sci. USA* 86:8402-8406).



## 24

Direct transformation techniques are suitable for any organism and cell type. In cases where DNA or RNA are injected or electroporated into plant cells, the plasmid used need not meet any particular requirements. Simple plasmids such as those from the pUC series may be used. If intact plants are to be regenerated from the transformed cells, it is necessary for an additional selectable marker gene to be present on the plasmid.

Stably transformed cells, i.e. those which contain the inserted DNA integrated into the DNA of the host cell, can be selected from untransformed cells when a selectable marker is part of the inserted DNA. By way of example, any gene which is capable of conferring resistance to antibiotics or herbicides (such as kanamycin, G 418, bleomycin, hygromycin or phosphinothricin and the like) is capable of acting as marker (see above). Transformed cells which express such a marker gene are capable of surviving in the presence of concentrations of such an antibiotic or herbicide which kill an untransformed wild type. Examples are mentioned above and preferably comprise the bar gene, which confers resistance to the herbicide phosphinothricin (Rathore KS et al. (1993) Plant Mol Biol 21(5):871-884), the nptII gene, which confers resistance to kanamycin, the hpt gene, which confers resistance to hygromycin, or the EPSP gene, which confers resistance to the herbicide Glyphosate. The selection marker permits selection of transformed cells from untransformed cells (McCormick et al. (1986) Plant Cell Reports 5:81-84). The plants obtained can be bred and hybridized in the customary manner. Two or more generations should be grown in order to ensure that the genomic integration is stable and hereditary.

30

The above-described methods are described, for example, in Jenes B et al. (1993) Techniques for Gene Transfer, in: Transgenic Plants, Vol. 1, Engineering and Utilization, edited by SD Kung and R Wu, Academic Press, pp.128-143, and in Potrykus (1991) Annu Rev Plant Physiol Plant Molec Biol 42:205-225). The construct to be expressed is preferably cloned into a vector which is suitable for transforming Agrobacterium tumefaciens, for example pBin19 (Bevan et al. (1984) Nucl Acids Res 12:8711f).

Once a transformed plant cell has been generated, an intact plant can be obtained using methods known to the skilled worker. For example, callus cultures are used as starting material. The development of shoot and root can be induced in this as yet undifferentiated cell biomass in the known fashion. The plantlets obtained can be planted out and used for breeding.



25

The skilled worker is familiar with such methods for regenerating plant parts and intact plants from plant cells. Methods which can be used for this purpose are, for example, those described by Fennell et al. (1992) Plant Cell Rep. 11: 567-570; Stoeger et al. (1995) Plant Cell Rep. 14:273-278; Jahne et al. (1994) Theor Appl Genet 89:525-533.

"Transgenic", for example in the case of a yeast TEP, refers to a nucleic acid sequence, an expression cassette or a vector comprising said TEP nucleic acid sequence or to an organism transformed with said nucleic acid sequence, expression cassette or vector or all those constructs established by recombinant methods in which either

- 15 a) the nucleic acid sequence encoding a yeast TEP or
- b) a genetic control sequence, for example a promoter which is functional in plant organisms, which is linked operably with said nucleic acid sequence under a)
- 20 c) (a) or (b)

are not in their natural genetic environment or have been modified by recombinant methods, it being possible for the modification to be, for example, a substitution, addition, deletion, inversion or insertion of one or more nucleotide residues. Natural genetic environment refers to the natural chromosomal locus in the source organism or the presence in a genomic library. In the case of a genomic library, the natural genetic environment of the nucleic acid sequence is preferably retained, at least to some extent. The environment flanks the nucleic acid sequence at least on one side and has a sequence length of at least 50 bp, preferably at least 500 bp, particularly preferably at least 1000 bp, very particularly preferably at least 5000 bp. A naturally occurring expression cassette, for example the naturally occurring combination of the promoter of a gene encoding for a yeast TEP with the corresponding yeast TEP gene, becomes a transgenic expression cassette when the latter is modified by non-natural, synthetic ("artificial") methods such as, for example, a mutagenization. Such methods are described (US 5,565,350; WO 00/15815; see also above).

Host or starting organisms which are preferred as transgenic organisms are, above all, plants in accordance with the above definition. Included for the purposes of the invention are all genera and species of higher and lower plants of the Plant



26

Kingdom, in particular plants which are used for obtaining oils, such as, for example, oilseed rape, sunflower, sesame, safflower, olive tree, soya, maize, wheat and nut species. Furthermore included are the mature plants, seed, shoots and seedlings, 5 and parts, propagation material and cultures, for example cell cultures, derived therefrom. Mature plants refers to plants at any desired developmental stage beyond the seedling stage. Seedling refers to a young, immature plant at an early developmental stage.

10

The transgenic organisms can be generated with the above-described methods for the transformation or transfection of organisms.

- 15 The invention furthermore relates to the use of the transgenic organisms according to the invention and to the cells, cell cultures, parts - such as, for example, in the case of transgenic plant organisms roots, leaves and the like - and transgenic propagation material such as seeds or fruits which are derived 20 therefrom for the production of foodstuffs or feedstuffs, pharmaceuticals or fine chemicals, in particular oils, fats, fatty acids or derivatives of these.

- Besides influencing the oil content, the transgenic expression 25 of a yeast TEP SEQ ID No: 1 or derivatives thereof in plants may mediate yet further advantageous effects such as, for example, an increased stress resistance. Such osmotic stress occurs for example in saline soils and water and is an increasing problem in agriculture. Increased stress tolerance makes it possible, 30 for example, to use areas in which conventional arable plants are not capable of thriving for agricultural usage.

- The invention now having been generally described will be more readily understood by reference to the following examples, which 35 are included for the purpose of illustration only, and are not intended to limit scope of the present invention.

#### Examples

#### 40 General methods:

Unless otherwise specified, all chemicals were from Fluka (Buchs), Merck (Darmstadt), Roth (Karlsruhe), Serva (Heidelberg) and Sigma (Deisenhofen). Restriction enzymes, DNA-modifying enzymes and molecular biological kits were from Amersham- 45 Pharmacia (Freiburg), Biometra (Göttingen), Roche (Mannheim), New England Biolabs (Schwalbach), Novagen (Madison, Wisconsin).



27

USA), Perkin-Elmer (Weiterstadt), Qiagen (Hilden), Stratagen (Amsterdam, Netherlands), Invitrogen (Karlsruhe) and Ambion (Cambridgeshire, United Kingdom). The reagents used were employed in accordance with the manufacturer's instructions.

5

For example, oligonucleotides can be synthesized chemically in the known manner using the phosphoamidite method (Voet, Voet, 2<sup>nd</sup> edition, Wiley Press New York, pages 896-897). The cloning steps carried out for the purposes of the present invention

10 such as, for example, restriction cleavages, agarose gel electrophoreses, purification of DNA fragments, transfer of nucleic acids to nitrocellulose and nylon membranes, linking DNA fragments, transformation of *E. coli* cells, bacterial cultures, multiplication of phages and sequence analysis of recombinant  
 15 DNA, are carried out as described by Sambrook et al. (1989) Cold Spring Harbor Laboratory Press; ISBN 0-87969-309-6. Recombinant DNA molecules were sequenced using an ABI laser fluorescence DNA sequencer following the method of Sanger (Sanger et al. (1977) Proc Natl Acad Sci USA 74:5463-5467).

20

#### EXAMPLE 1

Reduction of triacylglycerol accumulation in yeast cells lacking the *YJR098c* gene

25

Yeast strains used in this study were congenic to the W303-1A (Thomas & Rothstein, 1989) background. An *YJR098c* mutant strain, H1223, with the genotype *MATa yjr098c::HIS3 ADE2 can 1-100 his3-11,15 leu2-3, 112, trp1-1 ura3-1*, was generated as

30 described in Sandager et al., 2002. As a wild type control, we used the strain SCY62 *MATa ADE2 can 1-100 his3-11,15 leu2-3, 112 trp1-1 ura3-1*).

Yeast cells were cultivated at 30°C on a rotary shaker in liquid  
 35 synthetic medium (Sherman et al., 1986) supplemented with 2 % (wt/vol) glucose.

The lipid content of the yeast cells was determined as described by Dahlqvist et al. (2000) and is presented as nmol of fatty acid  
 40 (FA) per mg dry weight yeast.

The lipid content of a mutant yeast strain H1223, in which the *YJR098c* gene was disrupted, was analyzed and compared to wild type yeast cells (strain SCY62). The lipid content was determined  
 45 in yeast cells harvested in stationary phase after 50 hours of cultivation in liquid synthetic medium at 30°C. Lipids were extracted in chloroform, fractionated on TLC and quantified by



GC analyses (Dahlqvist et al., 2000). The total lipid content, measured as nmol fatty acids (FA) per dry weight yeast, in the YJR098c mutant yeast was 18% less than in the wild type, see table 1. The main reason for this difference was a lowered TAG content in the YJR098c mutant. Thus, the triacylglycerol amount in the mutant yeast was almost 36 % lower than in the wild type, whereas the polar lipid content only differed slightly between the YJR098 mutant and the wild type yeast, see table 1.

10 In summary, this experiment shows that the product of the YJR098c gene contributes to TAG accumulation in yeast.

Table 1. Lipid content in yeast disrupted in the YJR098c gene.

	control yeast (nmol FA/mg)	YJR098c - mutant (nmol FA/mg)
Sterol esters	28	25
Triacylglycerol	180	116
Other neutral lipids	7	9
Polar lipids	95	104
Total lipids	311	255

#### EXAMPLE 2

25 Increased accumulation of triacylglycerol in yeast cells expressing the YJR098c gene in combination with a strong promoter.

For induced high-level expression of the YJR098c gene, a 2439-bp DNA fragment, containing 29 bp up stream and 442 bp down stream of the gene, was amplified from wt W303 genomic DNA by using a 1:1 mixture of Taq and pfu DNA polymerases with the 5' primer, CTTGTAGAGGTTAACTGGGGA, and the 3' primer, TGAATTGTCCTCGCTGTCAA. The resulting PCR product was blunt end cloned into the BamHI site of the GAL1 yeast expression plasmid pUS10, which is a selection marker variant of the GAL1 yeast expression plasmid pJN92 (Ronne et al., 1991) thus generating the plasmid pUS30. pUS 10 was generated by removing the URA3 selection marker from the pJN92 plasmid by HindIII digestion and replacing it with the HIS3 gene, a 1768 bp DNA fragment that was blunt end cloned into the remaining part of the HINDIII digested pJN92. The wild type yeast strain SCY62 (MATa ADE2 can 1-100 his3-11,15 leu2-3, 112 trp1-1 ura3-1), was transformed with the pUS30 and cultivated at 28°C on a rotary shaker in synthetic medium (Sherman et al., 1986) lacking uracil and supplemented with 2 % (vol/vol) glycerol and 2 % (vol/vol) ethanol. The GAL1 promoter was induced after 6 or 24 hours of growth by the addition of 2 % (wt/vol) final



## 29

concentration of galactose. Cells were harvested after an additional 24 hours of growth. Wild type cells SCY62 (MATa ADE2 can 1-100 his3-11,15 leu2-3, 112 trp1-1 ura3-1) transformed with the empty vector, pUS10, and cultivated under identical conditions were used as a control. The lipid content of the yeast cells was determined as described by Dahlqvist et al. (2000) and is presented as nmol of fatty acid (FA) per mg dry weight yeast.

The effect of high-level expression of the YJR098c gene on lipid accumulation was studied by transforming the wild-type yeast strain SCY62 (Dahlqvist, et al., 2000) with a plasmid containing the YJR098c gene under control of the galactose-induced GAL1 promoter, see Table 2. High-level expression of the YJR098c gene from this promoter had no strong effect on the growth rate as determined by optical density measurements. The expression of the YJR098c gene was induced after 6 h (Table 2A) or 24 h (Table 2B) and cells were harvested after an additional 24 hours of cultivation. The total lipid content, determined as nmol fatty acids (FA) per mg yeast (Dahlqvist et al., 2000) in cells expressing the YJR098c gene from the GAL1 promoter was higher both at an early (Table 2A) or late (Table 2B) stationary growth stage as compared to cells transformed with an empty vector. The elevated lipid content in cells expressing the YJR098c gene from the GAL1 promoter was entirely explained by an increased TAG content whereas the content of polar lipids and sterol esters were unaffected.

In summary, the TAG content in yeast cells expressing YJR098c in combination with a strong promoter was increased with 26 to 28% as compared to the control (Table 2A and 2 B), which demonstrates the potential of the use of the YJR098c gene for increasing the oil content in transgenic organisms including yeast.

35

40

45



Table 2: Lipid content in yeast that expresses the YJR098c gene in combination with the GAL1 promoter

5	A	control yeast (nmol FA/mg)	High level of YJR098c expression (nmol FA/mg)
	Sterol esters	13	13
	Triacylglycerol	78	98
	Other neutral lipids	9	9
	Polar lipids	60	60
10	Total lipids	160	180
15	B	control yeast (nmol FA/mg)	High level of YJR098c expression (nmol FA/mg)
	Sterol esters	15	17
	Triacylglycerol	142	182
	Other neutral lipids	9	11
	Polar lipids	55	50
	Total lipids	221	260

## EXAMPLE 3

20

## Transgenic plants expressing YJR098c

For induced high level expression of the YJR098c gene in plants, a PCR fragment (2409 bp) was generated by the 5' primer (CTT GTA GAG GTT AAC TGG GGA) and the 3' primer (TGA ATT GTC CTC GCT GTC AA) adding 29 bases upstream of the gene and 442 bases downstream of the gene. The gene was cloned into the *Sma*I site of the vector pUC119, thus generating pUS 29. For *Agrobacterium*-mediated plant transformation a binary vector system including the primary cloning vector pART7 with a CaMV35S promoter and a binary pART27 vector (Gleave A., 1992) were used. The pART7 vector with a napin promoter is a construct where the napin promoter fragment (1101bp) described by Stålberg (1993) replaced the CaMV35S promoter from pART7 only loosing the *Xho*I site of the polylinker in the process. The YJR098c fragment were cut out from pUS 29 at the *Xba*I and *Sac*I site and then blunted into the pART7 vector with either the CaMV35S promoter, generating pEW 17 or with the napin promoter, generating pEW 14. The entire cartridge including the promoter, the YJR098c gene and a transcriptional termination region were removed from the pART7 vector as a *Not*I fragment and introduced directly to the pART27 vector. The plasmid was transformed into *Agrobacterium tumefaciens*.

Using floral dip essentially as described by Clough and Bent, 1998, plants of *Arabidopsis thaliana* were transformed with *Agrobacterium tumefaciens* GV3101 harboring either of the plasmids pEWART27-14 and pEWART27-17. Entire plants (inflorescence and ro-



## 31

sette) were submerged for 20 to 30 sec in the infiltration media consisting of 5% sucrose and 0.02% Silwet L-77 (Osi Specialties, Danbury, CT) plus resuspended transformed *A. tumefaciens* cells. Plants were then transferred to a growth chamber with a photoperiod of 16 h of light at 21°C and 8 h of dark at 18°C (70% humidity).

The seed oil content of T2 plants of the *Arabidopsis* transformants was analyzed by the use of conventional gas-liquid chromatography (GLC). As controls, seeds from wild type plants were used. The level of expression of the *YJR098c* gene in the seeds is determined by Northern blot analysis.

The result of the measurements for the lines comprising the *YJR098c* construct showed a significantly higher total oil content in transgenic lines compared to the measurements of wild-type plants.

## REFERENCES

- 20 Cases, S., Smith, S.J., Zheng, Y-W., Myers, H.M., Lear, S.R., Sande E., Novak, S., Collins, C., Welch, C.B., Lusi, A.J., Erickson, S.K., and Farese, R.V. (1998) *Proc. Natl. Acad. Sci. USA* 95, 13018-13023.
- 25 Dahlqvist, A., Stahl, U., Lenman, M., Banas, A., Lee, M., Sandager, L., Ronne, H. and Stymne, S. (2000) *Proc. Natl. Acad. Sci., USA* 97, 6487-6492.
- 30 Gleave, A. (1992) *Plant Molecular Biology* 20, 1203-1207.
- Lardizabal, K. D., Hawkins, D. J. and Thompson, G. A. (2001) *DGAT2: A New Diacylglycerol Acyltransferase Gene Family*. *JBC* 276 (42) 38862-38869.
- 35 Sandager, L., Gustavsson, M., Stahl, U., Dahlqvist, A., Wiberg, E., Banas, A., Lenman, M., Ronne, H., and Stymne, S. (2002) Storage lipid synthesis is non-essential in yeast. *Journal of Biological Chemistry* 277, 6478-6482
- 40 Sherman, F., Fink, G.R., and Hicks, J.B. (1986) *Laboratory Course Manual for Methods in Yeast Genetics*, Cold Spring Harbor Lab. Press, Plainview, NY.
- 45 Stålberg, K., Ellerström, M., Josefsson, L.-G., and Rask, L. (1993) *Plant Molecular Biology* 23, 671-683.



BASF Plant Science GmbH 20020437

Empf.zeit: 10/07/2002 17:08  
PF 53702 EP

32

Ronne, H., Carlberg, M., Hu, G.-Z. and Nehlin, J.O. (1991) Mol.  
Cell. Biol. 11, 4876-5884.

Thomas, B.J. and Rothstein, R. (1989) Cell 56, 619-630.

5

10

15

20

25

30

35

40

45



We claim:

1. A method of increasing the total oil content in a plant  
5 organism or a tissue, organ, part, cell or propagation  
material thereof, comprising
  - a) the transgenic expression of a polypeptide SEQ-ID No: 2  
10 from yeast in said plant organism or in a tissue, organ,  
part, cell or propagation material thereof, and
  - b) the selection of plant organisms in which - in contrast  
15 to or comparison with the starting organism - the total  
oil content in said plant organism or in a tissue, organ,  
part, cell or propagation material thereof is increased.
2. A method as claimed in claim 1, wherein the polypeptide from  
20 yeast is described by
  - a) a sequence with the SEQ ID NO: 2
  - b) a functional equivalent of a) with an identity of at  
least 60% of a sequence with SEQ ID NO: 2.
- 25 3. A method as claimed in claim 1 or 2, wherein the plant is  
an oil crop.
4. A method as claimed in claim 1 or 2, wherein the total oil  
30 content in the seed of a plant is increased.
5. A transgenic expression cassette comprising, under the  
control of a promoter which is functional in a plant organism  
or a tissue, organ, part, or cell thereof, a nucleic acid  
35 sequence SEQ ID No: 1.
6. A transgenic expression cassette as claimed in claim 5,  
wherein the nucleic acid sequence is described by
  - a) a sequence with the SEQ ID NO: 1  
40
  - b) a sequence derived from a sequence with the SEQ ID NO: 1,  
in accordance with the degeneracy of the genetic code
  - c) a sequence which has at least 60% identity with the  
45 sequence with the SEQ ID NO: 1.

Sequ.



34

7. A transgenic expression cassette as claimed in claim 5 or 6,  
wherein the promoter is a seed-specific promoter.
8. A transgenic vector comprising an expression cassette as  
5 claimed in any of claims 5 to 7.
9. A transgenic plant organism or tissue, organ, part, cell  
or propagation material thereof, comprising a polypeptide  
as defined in claim 2 or an expression cassette as claimed  
10 in any of claims 5 to 7 or a vector as claimed in claim 8.
10. A transgenic plant organism as claimed in claim 9, wherein  
the plant organism is selected from the group of the oil  
crops consisting of *Borvago officinalis*, *Brassica campestris*,  
15 *Brassica napus*, *Brassica rapa*, *Cannabis sativa*, *Carthamus*  
*tinctorius*, *Cocos nucifera*, *Crambe abyssinica*, *Cuphea*  
species, *Elaeis guinensis*, *Elaeis oleifera*, *Glycine max*,  
*Gossypium hirsutum*, *Gossypium barbadense*, *Gossypium*  
*herbaceum*, *Helianthus annuus*, *Linum usitatissimum*, *Oenothera*  
20 *biennis*, *Olea europaea*, *Oryza sativa*, *Ricinus communis*,  
*Sesamum indicum*, *Triticum* species, *Zea mays*, walnut and  
almond.
11. The use of a transgenic plant organism or tissue, organ,  
25 part, cell or propagation material thereof as claimed in  
claim 9 or 10 for the production of oils, fats, free fatty  
acids or derivatives of the above.

30

35

40

45



35

Use of a gene for increasing the oil content in plants

Abstract

5

The invention relates to methods for increasing the oil content in plants, preferably in plant seeds, by expressing a polypeptide from yeast. The invention furthermore relates to expression constructs for expressing the yeast polypeptide in plants, preferably in plant seeds, the transgenic plants expressing the yeast polypeptide and to the use of said transgenic plants for the production of food, feeds, seed, pharmaceuticals or fine chemicals, in particular for the production of oils.

15

20

25

30

35

40

45



## SEQUENCE LISTING

&lt;110&gt; BASF Plant Science GmbH

&lt;120&gt; Use of a gene for increase of oil content in plants

&lt;130&gt; 437-02

&lt;140&gt; 53702

&lt;141&gt; 2002-07-10

&lt;160&gt; 2

&lt;170&gt; PatentIn Ver. 2.1

&lt;210&gt; 1

&lt;211&gt; 2439

&lt;212&gt; DNA

&lt;213&gt; Saccharomyces cerevisiae

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (30) .. (1994)

&lt;400&gt; 1

```

ctt gtagagg ttaactgggg agtattaca atg atg gca act ccg gct act gat      53
                               Met Met Ala Thr Pro Ala Thr Asp
                               1                               5

ctt att tcc gat aat gat aaa tat aac aag caa tgt ctt tct gat tca      101
Leu Ile Ser Asp Asn Asp Lys Tyr Asn Lys Gln Cys Leu Ser Asp Ser
   10                               15                               20

agt gat agt ggt agt gat gta agc ttt ttt tcc gta aat gaa agc gaa      149
Ser Asp Ser Gly Ser Asp Val Ser Phe Phe Ser Val Asn Glu Ser Glu
   25                               30                               35                               40

ggt gaa ttg gat aca atg gag aaa gtg gat acc ttg att gga ggt gca      197
Gly Glu Leu Asp Thr Met Glu Lys Val Asp Thr Leu Ile Gly Gly Ala
                               45                               50                               55

aga gtt ata agc aat aaa gta gaa aaa gac agc gat agt gaa caa agg      245
Arg Val Ile Ser Asn Lys Val Glu Lys Asp Ser Asp Ser Glu Gln Arg
                               60                               65                               70

gga aga aag aag gaa aca act ggg ccc aat aac tat cat aat tta gaa      293
Gly Arg Lys Lys Glu Thr Thr Gly Pro Asn Asn Tyr His Asn Leu Glu
   75                               80                               85

gag aag caa gcg agt gcc att tct ctt gac gct gat gat gaa gat ctc      341
Glu Lys Gln Ala Ser Ala Ile Ser Leu Asp Ala Asp Asp Glu Asp Leu
   90                               95                               100

gat gaa att att tct tat tcc cat gac ggg aac tat gac agc tct cat      389
Asp Glu Ile Ile Ser Tyr Ser His Asp Gly Asn Tyr Asp Ser Ser His
  105                               110                               115                               120

```



2

aaa act ttc tcc ttt tcc tta cca ttt ggt aat aca aat ttc cga tca 437  
 Lys Thr Phe Ser Phe Ser Leu Pro Phe Gly Asn Thr Asn Phe Arg Ser  
 125 130 135

agt tca cca tta gcc ata att aaa act gtg ctt ccc aag act cca gat 485  
 Ser Ser Pro Leu Ala Ile Ile Lys Thr Val Leu Pro Lys Thr Pro Asp  
 140 145 150

gag ttc atc aaa aag aat cta aga aag aat gag atc aag caa aaa ctg 533  
 Glu Phe Ile Lys Lys Asn Leu Arg Lys Asn Glu Ile Lys Gln Lys Leu  
 155 160 165

aaa aaa tca acc tcc att tct tcc ttg gaa gag ata gaa tta ttt aaa 581  
 Lys Lys Ser Thr Ser Ile Ser Ser Leu Glu Glu Ile Glu Leu Phe Lys  
 170 175 180

tac gaa agg ggc att gat aat tca agg tta agg gct gtt aaa gaa tct 629  
 Tyr Glu Arg Gly Ile Asp Asn Ser Arg Leu Arg Ala Val Lys Glu Ser  
 185 190 195 200

ttg gaa atg gat gcc ttg aag aac tcc att aag caa ata aca gca gac 677  
 Leu Glu Met Asp Ala Leu Lys Asn Ser Ile Lys Gln Ile Thr Ala Asp  
 205 210 215

cca ttc gac aaa act cat gac gga tat tac cgt tcg cgt tta gaa tct 725  
 Pro Phe Asp Lys Thr His Asp Gly Tyr Tyr Arg Ser Arg Leu Glu Ser  
 220 225 230

ata tgg aat gaa ttg gaa gga gat gtc gtt ata atg ggt gga tat cga 773  
 Ile Trp Asn Glu Leu Glu Gly Asp Val Val Ile Met Gly Gly Tyr Arg  
 235 240 245

ggt agt gtg cta agg gat gct act act cat aag cga att tgg atc cca 821  
 Gly Ser Val Leu Arg Asp Ala Thr Thr His Lys Arg Ile Trp Ile Pro  
 250 255 260

tta aag gca ggt ttg aat atg acg aaa gtc gac tta ttg atc gga cct 869  
 Leu Lys Ala Gly Leu Asn Met Thr Lys Val Asp Leu Leu Ile Gly Pro  
 265 270 275 280

aat gac gaa gat gaa ctt aaa act cag aag gag att gtc cct gat gga 917  
 Asn Asp Glu Asp Glu Leu Lys Thr Gln Lys Glu Ile Val Pro Asp Gly  
 285 290 295

atg cta aca cat ata ggg cct gtt gat atc tct aag agg ttg ata aag 965  
 Met Leu Thr His Ile Gly Pro Val Asp Ile Ser Lys Arg Leu Ile Lys  
 300 305 310

agg cta gac gca aat cct aat tta aat gtt cag cag ttt ggc tat gat 1013  
 Arg Leu Asp Ala Asn Pro Asn Leu Asn Val Gln Gln Phe Gly Tyr Asp  
 315 320 325

tgg aga tta tcc ttg gac ata tct gcc aag cat tta acg act aaa cta 1061  
 Trp Arg Leu Ser Leu Asp Ile Ser Ala Lys His Leu Thr Thr Lys Leu  
 330 335 340

gag gaa att tac aat aag caa aaa aat aag aag gga ata tac atc att 1109



BASF Plant Science GmbH

20020437

PF 53702 BP

3

Glu Glu Ile Tyr Asn Lys Gln Lys Asn Lys Lys Gly Ile Tyr Ile Ile	
345	350 355 360
gcc cat tca atg ggc gga ttg gtc gca cat aaa gtg ttg caa gac tgt	1157
Ala His Ser Met Gly Gly Leu Val Ala His Lys Val Leu Gln Asp Cys	
	365 370 375
act cat ttg ata aga ggt att att tac gtg ggt tcc cca agc caa tgt	1205
Thr His Leu Ile Arg Gly Ile Ile Tyr Val Gly Ser Pro Ser Gln Cys	
	380 385 390
cca aat att tta ggt cct att agg ttt gga gat gat gtg atg tgg aat	1253
Pro Asn Ile Leu Gly Pro Ile Arg Phe Gly Asp Asp Val Met Trp Asn	
	395 400 405
aaa cta ttt tca cta aga acc aac ttt ttt atg aga agt agt ttc tat	1301
Lys Leu Phe Ser Leu Arg Thr Asn Phe Phe Met Arg Ser Ser Phe Tyr	
	410 415 420
ttt cta ccg tca gat ggt aga tgt ttt gtt gac aaa att acc tta gag	1349
Phe Leu Pro Leu Asp Gly Arg Cys Phe Val Asp Lys Ile Thr Leu Glu	
	425 430 435 440
agg tat gat ttc gat ttt ttt gat aca gat gtt tgg aaa acc ctt ggc	1397
Arg Tyr Asp Phe Asp Phe Phe Asp Thr Asp Val Trp Lys Thr Leu Gly	
	445 450 455
ttg tca cct ctc gtc aat gag aaa aga gag gaa tca gct cac gaa aaa	1445
Leu Ser Pro Leu Val Asn Glu Lys Arg Glu Glu Ser Ala His Glu Lys	
	460 465 470
tca aaa tta tta cca agg aaa acg aaa tca gcg ctt tcg ctt aaa gct	1493
Ser Lys Leu Leu Pro Arg Lys Thr Lys Ser Ala Leu Ser Leu Lys Ala	
	475 480 485
acc cta aac gca act acc aag ttt gtc cta aat gca cct gtt gtt agg	1541
Thr Leu Asn Ala Thr Thr Lys Phe Val Leu Asn Ala Pro Val Val Arg	
	490 495 500
aat gta gcc ggc aat aat aaa cag gta cca agg gat gtg cct ttc gat	1589
Asn Val Ala Gly Asn Asn Lys Gln Val Pro Arg Asp Val Pro Phe Asp	
	505 510 515 520
gaa gtc ttc cat aca tct tat gaa gat agc tgt gaa tat tta gcg aga	1637
Glu Val Phe His Thr Ser Tyr Glu Asp Ser Cys Glu Tyr Leu Ala Arg	
	525 530 535
act tta aaa cgt aca aag aat tat ttg gat agc tta gat tac gac ccg	1685
Thr Leu Lys Arg Thr Lys Asn Tyr Leu Asp Ser Leu Asp Tyr Asp Pro	
	540 545 550
aac aaa gaa cat cct cca ttg gcc atg gtt tac ggt aac aag gtt ccc	1733
Asn Lys Glu Tyr Pro Pro Leu Ala Met Val Tyr Gly Asn Lys Val Pro	
	555 560 565
act gtt aga ggt gct aaa gtg aac ggt ata caa gat ata aaa gat ggg	1781
Thr Val Arg Gly Ala Lys Val Asn Gly Ile Gln Asp Ile Lys Asp Gly	



**BASF Plant Science GmbH**

20020437

PF 53702 EF

4

570

575

580

aat tat gaa gat ttt tac tat ggt ccg ggc gac ggt gtt gtt cac cat 1829  
Asn Tyr Glu Asp Phe Tyr Tyr Gly Pro Gly Asp Gly Val Val His His  
585 590 595 600

aaa tgg tta ttg cct gaa cag aga ggc ttt cca gtt gtt tgt aaa atc 1877  
Lys Trp Leu Leu Pro Glu Gln Arg Gly Phe Pro Val Val Cys Lys Ile  
605 610 615

gcc agt tct tca ggt cat gtt agc tta atg acg gat ttg aaa tca atg 1925  
Ala Ser Ser Ser Gly His Val Ser Leu Met Thr Asp Leu Lys Ser Met  
620 625 630

gca aaa gca ttc ata tct atc gtc gac agc gaa aaa gaa gga aga aga 1973  
Ala Lys Ala Phe Ile Ser Ile Val Asp Ser Glu Lys Glu Gly Arg Arg  
635 640 645

tct cga aca cga act tct tca tgaaaggctt tttattcctt tgtttactat 2024  
Ser Arg Thr Arg Thr Ser Ser  
650 655

tcatatctgc atttttctttt traccaaaat tccgcctgct aaaaaaaatc tggcaacgca 2084

ccgcgaataa aaataaataa tattttttta tcttttagttg cctaaatact atttatttcg 2144

tcattttac aacctctttt atatacacca ttcgatttcc cacgaagtaa aataataatt 2204

ctataaacag atttatctga tatgctcaat ttccctccc attttcatta ttgtcccttc 2264

TGCTCTTCTT CGATGTCAAA ATTAACCTTC AGCCATAAGC TGCATGCGCT ACATTGGGGT 2324

aataattgat aaccagaatg actccgttcc atagcgtcta cattatcaat gcattcatct 2384

aacaaactct cactaaaatg aaaaccacca acaaattgac agcgaggaca attca 2439

<210> 2

<211> 655

<212> PRT

&lt;213&gt; Saccharomyces cerevisiae

<400> 2

Met Met Ala Thr Pro Ala Thr Asp Leu Ile Ser Asp Asn Asp Lys Tyr  
1 5 10 15

Asn Lys Gln Cys Leu Ser Asp Ser Ser Asp Ser Gly Ser Asp Val Ser  
 . 20 25 30

Phe Phe Ser Val Asn Glu Ser Glu Gly Glu Leu Asp Thr Met Glu Lys  
35 40 45

Val Asp Thr Leu Ile Gly Gly Ala Arg Val Ile Ser Asn Lys Val Glu  
50 55 60

Lys Asp Ser Asp Ser Glu Gln Arg Gly Arg Lys Lys Glu Thr Thr Gly  
65 70 75 80



Pro Asn Asn Tyr His Asn Leu Glu Glu Lys Gln Ala Ser Ala Ile Ser  
 85 90 95  
 Leu Asp Ala Asp Asp Glu Asp Leu Asp Glu Ile Ile Ser Tyr Ser His  
 100 105 110  
 Asp Gly Asn Tyr Asp Ser Ser His Lys Thr Phe Ser Phe Ser Leu Pro  
 115 120 125  
 Phe Gly Asn Thr Asn Phe Arg Ser Ser Ser Pro Leu Ala Ile Ile Lys  
 130 135 140  
 Thr Val Leu Pro Lys Thr Pro Asp Glu Phe Ile Lys Lys Asn Leu Arg  
 145 150 155 160  
 Lys Asn Glu Ile Lys Gln Lys Leu Lys Lys Ser Thr Ser Ile Ser Ser  
 165 170 175  
 Leu Glu Glu Ile Glu Leu Phe Lys Tyr Glu Arg Gly Ile Asp Asn Ser  
 180 185 190  
 Arg Leu Arg Ala Val Lys Glu Ser Leu Glu Met Asp Ala Leu Lys Asn  
 195 200 205  
 Ser Ile Lys Gln Ile Thr Ala Asp Pro Phe Asp Lys Thr His Asp Gly  
 210 215 220  
 Tyr Tyr Arg Ser Arg Leu Glu Ser Ile Trp Asn Glu Leu Glu Gly Asp  
 225 230 235 240  
 Val Val Ile Met Gly Gly Tyr Arg Gly Ser Val Leu Arg Asp Ala Thr  
 245 250 255  
 Thr His Lys Arg Ile Trp Ile Pro Leu Lys Ala Gly Leu Asn Met Thr  
 260 265 270  
 Lys Val Asp Leu Leu Ile Gly Pro Asn Asp Glu Asp Glu Leu Lys Thr  
 275 280 285  
 Gln Lys Glu Ile Val Pro Asp Gly Met Leu Thr His Ile Gly Pro Val  
 290 295 300  
 Asp Ile Ser Lys Arg Leu Ile Lys Arg Leu Asp Ala Asn Pro Asn Leu  
 305 310 315 320  
 Asn Val Gln Gln Phe Gly Tyr Asp Trp Arg Leu Ser Leu Asp Ile Ser  
 325 330 335  
 Ala Lys His Leu Thr Thr Lys Leu Glu Glu Ile Tyr Asn Lys Gln Lys  
 340 345 350  
 Asn Lys Lys Gly Ile Tyr Ile Ile Ala His Ser Met Gly Gly Leu Val  
 355 360 365  
 Ala His Lys Val Leu Gln Asp Cys Thr His Leu Ile Arg Gly Ile Ile  
 370 375 380



BASF Plant Science GmbH

20020437

PF 53704 EF

6

Tyr Val Gly Ser Pro Ser Gln Cys Pro Asn Ile Leu Gly Pro Ile Arg  
385 390 395 400

Phe Gly Asp Asp Val Met Trp Asn Lys Leu Phe Ser Leu Arg Thr Asn  
405 410 415

Phe Phe Met Arg Ser Ser Phe Tyr Phe Leu Pro Leu Asp Gly Arg Cys  
420 425 430

Phe Val Asp Lys Ile Thr Leu Glu Arg Tyr Asp Phe Asp Phe Phe Asp  
435 440 445

Thr Asp Val Trp Lys Thr Leu Gly Leu Ser Pro Leu Val Asn Glu Lys  
450 455 460

Arg Glu Glu Ser Ala His Glu Lys Ser Lys Leu Leu Pro Arg Lys Thr  
465 470 475 480

Lys Ser Ala Leu Ser Leu Lys Ala Thr Leu Asn Ala Thr Thr Lys Phe  
485 490 495

Val Leu Asn Ala Pro Val Val Arg Asn Val Ala Gly Asn Asn Lys Gln  
500 505 510

Val Pro Arg Asp Val Pro Phe Asp Glu Val Phe His Thr Ser Tyr Glu  
515 520 525

Asp Ser Cys Glu Tyr Leu Ala Arg Thr Leu Lys Arg Thr Lys Asn Tyr  
530 535 540

Leu Asp Ser Leu Asp Tyr Asp Pro Asn Lys Glu Tyr Pro Pro Leu Ala  
545 550 555 560

Met Val Tyr Gly Asn Lys Val Pro Thr Val Arg Gly Ala Lys Val Asn  
565 570 575

Gly Ile Gln Asp Ile Lys Asp Gly Asn Tyr Glu Asp Phe Tyr Tyr Gly  
580 585 590

Pro Gly Asp Gly Val Val His His Lys Trp Leu Leu Pro Glu Gln Arg  
595 600 605

Gly Phe Pro Val Val Cys Lys Ile Ala Ser Ser Ser Gly His Val Ser  
610 615 620

Leu Met Thr Asp Leu Lys Ser Met Ala Lys Ala Phe Ile Ser Ile Val  
625 630 635 640

Asp Ser Glu Lys Glu Gly Arg Arg Ser Arg Thr Arg Thr Ser Ser  
645 650 655



**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**